



Case Report

Journal Homepage: <http://crp.tums.ac.ir>

Tracheobronchopathia Osteochondroplastica

Ehsan Sanaei^{1*}, Fatemeh Aghaei Meybodi¹, Marzie Vaghefi²

1. Department of Internal Medicine, School of Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.
2. Department of Pediatrics, School of Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.



Citation Sanaei E, Aghaei Meybodi F, Vaghefi M. Tracheobronchopathia Osteochondroplastica. Case Reports in Clinical Practice. 2019; 4(1):23-26.

Running Title: Tracheobronchopathia Osteochondroplastica (TPO)

**Article info:****Received:** 17 January 2019**Revised:** 09 February 2019**Accepted:** 22 February 2019**Keywords:**Tracheobronchopathia
Osteochondroplastica; Breathlessness**ABSTRACT**

Tracheobronchopathia Osteochondroplastica (TPO) is a rare benign disorder of the lower part of the trachea and the upper part of the main bronchi. A 55-year-old female patient had been referring to our pulmonary and critical care center suffering from chronic dry cough and breathlessness on exertion and retrosternal pain for several months who was admitted to our center when she was complaining about breathlessness on exertion and chronic dry cough and retrosternal pain.

Introduction

Tracheobronchopathia Osteochondroplastica (TPO) is a rare benign disorder of the lower part of the trachea and the upper part of the main bronchi [1-3]. It was first described in the mid-19th century, and approximately 300 cases have been reported since then.

A higher incidence of TPO was seen in northern Europe countries, especially in Finland [4]. Because many cases are asymptomatic, TPO is mainly diagnosed post-mortem. Symptoms can range from productive or non-productive cough, hemoptysis, dyspnea, dryness of the throat, and recurrent pulmonary infections (e.g. retention pneumonia) [4].

*** Corresponding Author:****Ehsan Sanaei, MD. Candidate****Address:** Department of Internal Medicine, School of Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.**E-mail:** ehsan.bmsu@gmail.com




Figure 1. Increasing thickness of the trachea

In severe cases, the diagnosis is made during a difficult intubation. The characteristic bronchoscopic findings are reported as beaded, speculated, rock garden, cobble stoned, or stalactite grotto appearance. The typical histological appearance confirms the diagnosis.

Case Presentation

A 55-year-old female was referred to our pulmonary and critical care center suffering from chronic dry cough and breathlessness on exertion and retrosternal pain since several months ago. She was finally admitted to our center with the complaint of breathlessness on exertion, chronic dry cough, and retrosternal pain. However, her endoscopy and cardiac examination, echocardiography, and coronary artery angiography were normal. She was not a smoker with no history of allergy. Her lung auscultation, lung function tests, and other physical examination were normal. Regarding her laboratory report, her Erythrocyte Sedimentation Rate (ESR)




Figure 2. Increasing thickness of the trachea in the form of nodular and calcification

and C-Reactive Protein (CRP), Angiotensin-Converting Enzyme (ACE) level, cytoplasmic-ANCA and perinuclear-ANCA were within normal range. Increasing the thickness of the trachea in the chest radiography was seen (Figure 1).

In computed tomography of the chest, the increased thickness of the trachea in the form of nodular and calcification was seen (Figure 2 and 3). So, flexible fiberoptic bronchoscopy was done that revealed diffuse calcified nodularity of the trachea and main and lobar bronchus (Figure 4). Then, multiple biopsies were taken. Histological examination revealed bronchial cartilage and lamellar bone with little marrow (Figure 5). Smears and cultures for *Mycobacterium tuberculosis* were all negative.



Figure 3. Increasing thickness of the trachea in the form of nodular and calcification (lung view)



Figure 4. Diffuse calcified nodularity of the trachea




Figure 5. Bronchial cartilage with abnormal and unevenly distributed mineralization supporting the diagnosis of Tracheobronchopathia Osteochondroplastica

Discussion

In the bronchoscopic view, TPO appears as whitish, hard spicules projecting into the tracheal lumen from the anterior and lateral walls, sparing the posterior wall. Also, the larynx and the main bronchi could be involved. The diagnosis of TPO is confirmed by typical histological findings, usually from biopsies or postmortem analysis. Our patient suffered from exertional dyspnea and chronic dry cough.

Moreover, TPO nodules may also be caused by endobronchial sarcoidosis, calcification lesions of tuberculosis, papillomatosis, malignant lesions, and tracheobronchial calcinosis [5-10]. Some patients were initially thought to have asthma [11] or bronchial/trachea tumors.

Because typical symptoms are absent, TPO is most likely underdiagnosed. In this case, the patient was referred to the clinic with breathlessness on exertion, chronic dry cough and retrosternal pain while her endoscopy, cardiac examination, echocardiography, and coronary artery angiography reports were normal. Only severe cases suffer from wheezing and dyspnea caused by the obstruction of the airway lumen. Hemoptysis can occasionally result from the ulceration of a nodule or an acute infection. Sometimes TPO causes difficulties in endotracheal intubation [12, 13]. In most cases, the disease progresses very slowly although it may eventually end in respiratory insufficiency [14].

Because no specific therapy is available, treatment is only symptomatic, which includes antibiotics in case of

bacterial infections, mechanical measures to remove obstruction nodules by using cryotherapy, laser excision, external beam irradiation, radiotherapy, stent insertion, interventional bronchoscopy, or surgical resection [15, 16]. Also, successful recovery with nonsteroidal anti-inflammatory drugs and dapsone has been reported in symptomatic cases.

For our patient, the first treatment was prescription of ICS fluticasone, naproxen 500 mg q 12 h, and pregabalin 75 daily respiratory unite of Shahid Sadoughi University of Medical Sciences. After 4 weeks, because of unsuccessful treatment of the symptoms, the patient was referred to another center for interventional bronchoscopy.

Ethical Considerations

Compliance with ethical guidelines

There was no ethical considerations to be considered in this research.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflict of interest

The authors declared no conflict of interest.

References

- [1] Vilkmán S, Keistinen T. Tracheobronchopathia Osteochondroplastica: Report of a young man with severe disease and retrospective review of 18 cases. *Respiration*. 1995; 62(3):151-4. [DOI:10.1159/000196411]
- [2] Karlıkaya C, Yüksel M, Kiliçli S, Candan L. Tracheobronchopathia Osteochondroplastica. *Respirology*. 2000; 5(4):377-80. [DOI:10.1046/j.1440-1843.2000.00278.x]
- [3] Jabbarjarani HR, Radpey B, Kharabian S, Masjedi MR. Tracheobronchopathia Osteochondroplastica: Presentation of ten cases and review of the literature. *Lung*. 2008; 186(5):293. [DOI:10.1378/chest.134.4_MeetingAbstracts.p10002]
- [4] Prakash UB. Tracheobronchopathia Osteochondroplastica. *Seminars in Respiratory and Critical Care Medicine*. 2002; 23(2):167-75. [DOI:10.1055/s-2002-25305] [PMID]
- [5] Kart L, Kiraz K, Büyükoğlan H, Ozesmi M, Sentürk Z, Gülmez I, et al. Tracheobronchopathia Osteochondroplastica: Two cases and review of literature. *Tüberk Toraks*. 2004; 52(3):268-71. [PMID]

- [6] Neumann A, Kasper D, Schultz-Coulon HJ. [Clinical aspects of tracheopathia osteoplastica (German)]. *HNO*. 2001; 49(1):41-7. [\[DOI:10.1007/s001060050706\]](https://doi.org/10.1007/s001060050706)
- [7] Sutor GC, Glaab T, Eschenbruch C, Fabel H. Tracheobronchopathia Osteochondroplastica: An uncommon cause of retention pneumonia in an elderly patient. *Pneumologie*. 2001; 55(12):563-7. [\[DOI:10.1055/s-2001-19000\]](https://doi.org/10.1055/s-2001-19000)
- [8] Hantous-Zannad S, Sebai L, Zidi A, Khelil JB, Mestiri I, Besbes M, et al. Tracheobronchopathia Osteochondroplastica presenting as a respiratory insufficiency: Diagnosis by bronchoscopy and MRI. *European Journal of Radiology*. 2003; 45(2):113-6. [\[DOI:10.1016/S0720-048X\(02\)00028-1\]](https://doi.org/10.1016/S0720-048X(02)00028-1)
- [9] Meyer CN, Døssing M, Broholm H. Tracheobronchopathia Osteochondroplastica. *Respiratory Medicine*. 1997; 91(8):499-502. [\[DOI:10.1016/S0954-6111\(97\)90117-7\]](https://doi.org/10.1016/S0954-6111(97)90117-7)
- [10] Gillissen A, Richter A, Oster H. Clinical efficacy of short-term treatment with extra-fine HFA beclomethasone dipropionate in patients with post-infectious persistent cough. *Journal of Physiology and Pharmacology*. 2007; 58(5):223-32. [\[PMID\]](https://pubmed.ncbi.nlm.nih.gov/17111111/)
- [11] Restrepo S, Pandit M, Villamil MA, Rojas IC, Perez JM, Gascue A. Tracheobronchopathia Osteochondroplastica: Helical CT findings in 4 cases. *Journal of Thoracic Imaging*. 2004; 19(2):112-6. [\[DOI:10.1097/00005382-200404000-00010\]](https://doi.org/10.1097/00005382-200404000-00010)
- [12] Khan AM, Shim C, Simmons N, Chung V, Alterman DD, Haramati LB, et al. Tracheobronchopathia Osteochondroplastica: A rare cause of tracheal stenosis- "TPO stenosis". *The Journal of Thoracic and Cardiovascular Surgery*. 2006; 132(3):714-6. [\[DOI:10.1016/j.jtcvs.2006.05.024\]](https://doi.org/10.1016/j.jtcvs.2006.05.024)
- [13] Coetmeur D, Bovyn G, Leroux P, Niel-Duriez M. Tracheobronchopathia Osteochondroplastica presenting at the time of a difficult intubation. *Respiratory Medicine*. 1997; 91(8):496-8. [\[DOI:10.1016/S0954-6111\(97\)90116-5\]](https://doi.org/10.1016/S0954-6111(97)90116-5)
- [14] Molloy AR, McMahon JN. Rapid progression of tracheal stenosis associated with tracheopathia osteo-chondroplastica. *Intensive Care Medicine*. 1988; 15(1):60-2. [\[DOI:10.1007/BF00255642\]](https://doi.org/10.1007/BF00255642)
- [15] Shigematsu Y, Sugio K, Yasuda M, Sugaya M, Ono K, Takenoyama M, et al. Tracheobronchopathia Osteochondroplastica occurring in a subsegmental bronchus and causing obstructive pneumonia. *The Annals of Thoracic Surgery*. 2005; 80(5):1936-8. [\[DOI:10.1016/j.athoracsur.2004.06.101\]](https://doi.org/10.1016/j.athoracsur.2004.06.101)
- [16] Baugnee PE, Delaunois LM. Mycobacterium avium-intracellulare associated with Tracheobronchopathia Osteochondroplastica. *European Respiratory Journal*. 1995; 8(1):180-2. [\[DOI:10.1183/09031936.95.08010180\]](https://doi.org/10.1183/09031936.95.08010180)